EXHIBIT NO.

21

Watching Briefs

Report on the 2018 Acute Flaccid Myelitis Outbreaks in the USA

Author:

Ĭ

Abstract

An increase in the number of Acute Flaccid Myelitis (AFM) cases reported in the USA was detected in August 2018. The CDC has confirmed 186 cases of AFM in 2018 making this the largest outbreak of AFM reported in the USA. The cause however remains unknown. In this watching brief we review key features of the current outbreak in context, describe what is known about AFM, and consider probable causes. A two-year epidemic cycle of AFM appears to be emerging, and most current evidence implicates both EV-A71 and EV-D68 as the probable cause of the current outbreak yet key questions remain.

Keywords:

How to Cite: Adam, D.C., 2019. Report on the 2018 Acute Flaccid Myelitis

Outbreaks in the USA. Global Biosecurity, 1(1), pp.129–139.

DOI: Her de

Views

Downloads

Report on the 2018 Acute Flaccid Myelitis Outbreaks in the USA

Date of first report of

the outbreak

28-Aug-18

Disease or outbreak

Acute Flaccid Myelitis (AFM)

Origin (country, city,

region)

Multistate, USA

Suspected Source

(specify food source, Non-polio enteroviruses EV-A71 and EV-D68 (,)

zoonotic or human origin or other)

The exact date of outbreak onset is not publicly known. There have been sporadic cases of AFM reported in the US since February 2018, however data provided by the Centres for Disease Control and Prevention (CDC) suggests that an increase in reported AFM cases occurred in late August 2018 (,). Late August as a possible date of outbreak beginning is supported by an October 10th statement from the Washington State Department of Health describing an initial cluster of five AFM cases with symptom onset dated August 28, 2018 (). Supporting this, the CDC later confirmed that an increased number of reports for suspected AFM were received with symptom onset dates during August, September, and October ().

Date of outbreak beginning Date outbreak declared over

N/A - Ongoing as of 28 December 2018.

The CDC is reporting at least 38 US States with confirmed AFM cases reported since the beginning of 2018 (). Of the 341 suspected cases reported since February, 186 have been confirmed as of December 28th 2018 (). A comprehensive list of states affected with individual counts since February can be seen here: https://www.cdc.gov/acute-flaccid-myelitis/afm-surveillance.html

It remains difficult to determine whether state-wise totals reported by the CDC comprise sporadic cases reported prior to late August (the apparent onset of the epidemic wave) or as part of the epidemic. Confirmation of cases reported during the epidemic are also uninformative as large numbers of suspected

Affected countries & regions

cases could have onset dates between February and August, and individual case data are not described. Public reporting, however, implicates at least seven states in the epidemic wave beginning in late August which could have seeded other states previously reporting sporadic cases prior to August.

These include:

- 1. Washington (1) (10 confirmed)
- 2. Illinois () (9 confirmed)
- 3. Colorado () (16 confirmed)
- 4. Minnesota (1) (10 confirmed)
- 5. New Jersey (1) (10 confirmed)
- 6. Pennsylvania () (9 confirmed)
- 7. Ohio () (12 confirmed)

AFM reported since the beginning of 2018. Only 186 cases have been clinically confirmed, with an unknown number still under investigation. At least half of these confirmed cases have been reported between late August and November 2018 (,). AFM is a clinical syndrome that principally includes a sudden onset of acute limb weakness or paralysis (). Other features of AFM include difficulty breathing, ataxia (unsteady walking), wobbliness, headache, stiff neck, dizziness and jerking movements (). Fever can also be expected in 50% of cases ().

As of December 28, there have been 341 suspected cases of

Number of cases (specify at what date if ongoing)

Clinical features

flaccid limb weakness" and "evidence of spinal cord lesions" within the grey matter as shown by Magnetic Resonance Imaging (MRI) (). Evidence of increased white blood cell

Confirmation of AFM requires clinical evidence of "acute

count within the Cerebrospinal fluid (CSF) may also be supportive, but not confirmatory, of AFM (). In the current outbreak, the CDC confirmed that 78% of cases confirmed prior to November 2nd (N=62/80) had respiratory symptoms such as cough, runny nose and congestion, 81% (N=65/80) had fever, and 38% (N=30/80) has gastrointestinal symptoms ().

AFM as a clinical syndrome may be caused by different pathogens and cannot itself be transmitted from person-to-person. Infectious causes of AFM include polio virus, non-polio enteroviruses EV-A71 and EV-D68 (), some flaviviruses, adenoviruses type 21, and some herpesviruses including cytomegalovirus and Epstein-barr virus (, ,). Non-infectious causes may include environmental toxins or genetic disorders ().

Transmission of EV-A71 is by the faecal-oral route, however the virus is also present and can transmit via respiratory secretions such as nasal mucus ('). EV-D68 has only recently been associated with AFM (,), and is primarily transmitted via the respiratory route and respiratory illness is the main clinical syndrome for EV-D68 infection (,). EV-D68 has been detected in faeces of some infected cases but faecal-oral route transmission has not been established like EV-A71 (). Both EV viruses can survive on surfaces as fomites (,).

Mode of transmission (,).
(dominant mode and other documented modes)

Other kn

Other known causes of AFM include West Nile, Saint Louis encephalitis, and Japanese encephalitis viruses, which are all

mosquito-borne flaviviruses and are primarily carried by Culex spp. mosquitoes (14).

Adenovirus transmission is similar to EV-D68, primarily via respiratory secretions. However, unlike EV-D68, transmission via the faecal-oral route has been shown (___).

The exact demographics of all cases in this outbreak are not publicly known. The CDC has reported that among cases confirmed prior to November 2nd, 59% (47/80) were male (). The median age of cases is 4 years (). This is consistent with the consensus among reporting media indicating that 90% of cases affected are aged under 18 years (). In the August-November epidemic wave, all cases in Washington are aged under six years (), all cases in Minnesota are under 10 (), and all cases in Illinois are under the age of 18 (). In Colorado, Pennsylvania, and Ohio, all cases are described among "children" (,,,). There are no details on the cases in New

Demographics of cases

Jersey yet.

There has been one suspected death from AFM diagnosed in May 2018, however the CDC has yet to confirm this (). In the current epidemic wave, starting in August, no fatalities have been confirmed or reported.

Case fatality rate

Mild cases of AFM can expect to fully recover, however at least half of cases may have residual limb weakness requiring ongoing physical therapy (). Severe cases of AFM may experience respiratory failure and require ventilator support (). Rare neurological complications associated with AFM can sometimes lead to death (). Long-term complications include paralysis and residual limb deficits.

Complications

The causes of AFM are varied and include both infectious and non-infectious aetiologies. As such, the CDC recommends a variety of prevention strategies including: staying up-to-date with vaccines, basic hygiene i.e. washing your hands, and protecting yourself from mosquito bites (). Of the known causes of AFM, there are no commercially available vaccines for EV-A71 (), EV-D68 (), Saint Louis Encephalitis virus () or West Nile virus () in the US. A vaccine is available to prevent Japanese Encephalitis virus, however it is only recommended for travellers visiting endemic countries for longer than one-month and not available for infants under two months of age ().

Available prevention

There is no available treatment for AFM, which is limited to supportive care only. This includes the use of assisted ventilation when necessary (). Specialists may assist with physical therapy whilst hospitalised to reduce muscle weakness or loss (). The antiviral Fluoxetine has been trailed as treatment for presumptive EV-D68 associated AFM. However,

Available treatment

it was not associated with improved outcomes (). In recent times, large outbreaks of AFM were first documented in the US in 2014. In the 2014, 120 cases of AFM were confirmed across 34 states between August and December in the US (). This AFM outbreak was temporally and geographically coincident with large clusters of severe EV-D68 infections. A second outbreak of AFM was reported two years later in 2016 in the US, with 149 confirmed cases across 39 states including the District of Columbia, again coincident with reports of large EV-D68 detections. Application of the

Comparison with past outbreaks

Bradford-Hill criteria concluded for the first time that EV-D68 was the likely the cause of the 2014 AFM outbreak (, , ,). In each intervening year, in 2015 and 2017, sporadic cases of AFM have been reported: 22 confirmed cases across 17 states. and 33 confirmed cases across 16 states respectively. Unlike previous AFM outbreaks associated with EV-D68, this outbreak has been associated with both EV-A71 and EV-D68. Among 125 confirmed AFM cases tested, 21 (42%) have tested positive for EV-A71, 16 (32%) positive for EV-D68, and 13 (26%) positive for other viruses types included rhinovirus A subtypes and Coxsackie A viruses (). In some states, one species appears to predominate, such as in Colorado, where among 16 AFM cases, 11 tested positive for EV-A71, one for EV-D68, and two negative for enterovirus species (). Since 2014, there have been 512 confirmed cases of AFM in the US (). To-date, this is the largest outbreak of confirmed AFM in the US. The incidence of reported AFM has increased every two years since 2014 in the United States. Surveillance data shows 120 cases of confirmed AFM reported in 2014, 149 confirmed in 2016, and 193 in 2018 confirmed so far (,). The number of states implicated in each outbreak has also increased from 34 in 2014 to 39 in both 2016/18 (). Reporting of AFM however remains voluntary in the US so the true incidence remains unknown. The clinical similarity of AFM to other neurologic syndromes, e.g. Guillain-Barre syndrome and idiopathic transverse myelitis, increases the chance of

misdiagnosis, potentially concealing the true AFM incidence

further (). Interestingly, the onset of the 2018 outbreak

Unusual features

Critical analysis

coincides with time-of-onset observed in both previous outbreaks in 2014 and 2016: August to October (). This suggests a two-year cyclical dynamic of AFM outbreaks and a seasonal pattern is emerging, meaning another outbreak in August 2020 might also be expected. This cyclical dynamic also supports the role of a viral agent as the cause of the 2018 outbreak rather than unknown sporadic or environmental cause, such as a toxin. Supporting this conclusion, the CDC released a report on 16 November 2018 based on the results of early diagnostic testing stating "clinical, laboratory, and epidemiologic evidence to date suggest a viral association" (). Among 125 confirmed AFM cases tested, 21 (42%) have tested positive for EV-A71, 16 (32%) positive for EV-D68, and 13 (26%) positive for other viruses types, including rhinovirus A subtypes and Coxsackie A viruses (). Detection in the cerebrospinal fluid (CSF) is confirmative of the pathogen being the cause of AFM, however only two AFM cases among 21 specimens tested positive: one EV-A71, and another for EV-D68. While these numbers are limited, it is rare that EV-D68 and EV-A71 are detected in the CSF, even among cases with confirmed AFM and positive respiratory or stool samples. For example, in the 2014 outbreak, of the 120 confirmed AFM cases, only 12 had confirmed EV-D68 infections, 11 from respiratory samples, and one from CSF (,). This is likely due to the limited replication and persistence of EV virions in the CSF (, ,) and the delay between symptom onset and specimen sampling as observed in the 2014 outbreak () .Therefore, while CSF positive detections are confirmatory,

negative CSF results does not weaken other clinical and diagnostic evidence for EV-A71 and EV-D68 aetiology. Due to the emergence of EV-D68 as a likely cause of the 2014/16 AFM outbreak in US, it is possible that EV-D68 may be the cause of the 2018 outbreak. Clinical and diagnostic evidence provided by the CDC above supports this conclusion. Perhaps unusual in this outbreak is the simultaneous detection of EV-A71 among many AFM cases. Detection of EV-A71 in previous AFM outbreaks in the US is not known. In 2014, 11 Non-EV-D68 enterovirus species were detected in the stool of AFM cases, suggestive of EV-A71 infection as a faecal-oral pathogen but was not specified (). In 2016, a single AFM case in Washington State tested positive for EV-A71 among 10 confirmed AFM cases (). Similarly, there is little evidence internationally for co-circulation of EV-D68 and EV-A71 among AFM cases. For example, in France, enterovirusassociated AFM appears mostly associated with EV-A71, although EV-D68 has been detected previously in a single case). Of all the 38 states affected, Texas has recorded the most AFM cases (n=25), followed by Colorado (n=16). Based on public data and reporting, we cannot determine if the majority of confirmed cases in Texas had onset dates during the epidemic period August to October, or prior. In September, a Morbidity and Mortality Weekly Report (MMWR) was published describing an increase in EV-A71 associated encephalitis, meningitis and AFM in Colorado between May 10 and June 5, 2018 (). During this period, 34 young children (<2 years) presented with neurological symptoms and tested

positive for EV-A71; three were subsequently classified as AFM (). In the current epidemic which began in August, of the 16 positive AFM cases, 11 have tested positive for EV-A71 (). Only one cases has tested positive one for EV-D68, while two have tested negative for enterovirus species (). This might suggest EV-A71 is the predominate cause of the current AFM outbreak in Colorado, while EV-D68 is only a minor cause there.

In 2014, like 2018, Colorado was also one of the first states to report an increase in AFM detections (). This suggests Colorado may be a significant state for the emergence of AFM epidemics in the US. It is unclear if other states also experience simultaneous emergence of AFM or if a single state such as Colorado may act as a source, potentially disseminating the disease across the US. If the emergence is simultaneous, the environment and seasonality of non-polio enteroviruses may provide one explanation. Non-polio enteroviruses are known to circulate all-year round, however peaks typically occur in summer between July and September in the Northern Hemisphere (). Outbreak timing and transmission intensity has been shown to be associated with latitudes and dew-point temperatures such that northern states observed peaks later in the season whereas southern states observed a regular distribution of cases throughout the year. This supports the case for non-polio enteroviruses as the cause of the current AFM outbreak as most states affected since August epidemic wave are located in the Northern US: Washington, Illinois, Colorado, Minnesota, New Jersey, Pennsylvania, and Ohio (

of EV-D68 have also been reported in New York, however none of these cases have exhibited symptoms associated with AFM (____). Other evidence suggests non-polio enterovirus seasonality may be driven by a waning of seroprevalence in the population and the emergence of immunologically naïve newborns (____).

The detection of 13 non-enterovirus species among AFM cases () means other known viral aetiologies might also be implicated, including rhinovirus A subtypes and Coxsackie A viruses, however they do not have a history of neurological invasion and AFM. Likewise, symptomatic evidence can exclude flaviviruses previously associated with AFM such as West Nile Virus, which typically present with skin rash, as the cause of this AFM outbreak. Furthermore, diagnostic testing for flaviviruses require blood samples, which if taken, have not been publicly reported. Adenoviruses type 21 (AD21) has also been associated with AFM in the past (). A large outbreak of Adenovirus has sickened 36 in New Jersey between September and November, which has temporal and spatial associations with current AFM outbreaks also in New Jersey (;,). Eleven deaths have been implicated in this Adenovirus outbreak, however no cases have developed AFM and the dominant clinical syndrome was reported to be respiratory illness (). Typing has also shown a mix of adenoviruses types 3 and 7 predominating in the New Jersey outbreak, which do not have a history of causing AFM (). This might practically eliminate AD21 as a potential cause of the current AFM outbreak. What is the cause of AFM in the US in 2018?

Key questions

Why is AFM increasing in the US?

Is Colorado a key state for the emergence of AFM epidemics in the US or is the seasonality of non-polio enteroviruses a key predictor?

Is EV-D68 or EV-A71 the primary cause of this AFM epidemic, and if so, how common is it for two different co-circulating viruses to cause a single outbreak?

Is there common factor among all cases with positive and

Is there common factor among all cases with positive and negative enterovirus results?

Is there another non-enterovirus related cause to this outbreak? Could Rhinovirus A and Coxsackie A viruses be a new cause of AFM?

Can we exclude flaviviruses (West Nile, Saint Louis encephalitis virus, and Japanese encephalitis virus), other adenoviruses or a new pathogen?

References

1. McKay SL. Increase in Acute Flaccid Myelitis—United States, 2018. MMWR Morbidity and mortality weekly report. 2018;67.

DOI:

- 2. Stelzer-Braid S, Rawlinson W. Outbreaks of acute flaccid myelitis in the US. BMJ. 2018;363:k5246. DOI:
- 3. CDC. AFM Confirmed U.S. Cases [updated 7 January 2019. Available from:
- 4. CDC. AFM Investigation 2018 [updated 2 January 2019. Available from:
- 5. WSDH. Investigation into five possible AFM illnesses from four WA counties 2018 [updated 10 October 2018. Available

from:

6 IDPH Statement on Acute Flaccid Myelitis 2018 Jundated 10 October 2018, Avoil

6. IDPH. Statement on Acute Flaccid Myelitis 2018 [updated 10 October 2018. Available from: The Company of the Market and the Company of the

7. CDPHE. Statement from CDPHE on enterovirus and acute flaccid myelitis 2018 [updated 9 October 2018. Available

from: The second of the second second

8. MDH. Statement on cases of acute flaccid myelitis 2018 [updated 5 October 2018. Available from:

- 9. NJDOH. Acute flaccid myelitis (AFM) 2018 [updated 5 November 2018. Available from:
- 10. PDOH. Acute Flaccid Myelitis 2018 [updated 7 November 2018. Available from:
- 11. Cass A. Polio-like virus AFM remains rare, but cases increasing in Ohio: The News-Herald; 2018 [updated 8 December 2018. Available from:
- 12. CDC. About Acute Flaccid Myelitis 2018 [updated 22 October 2018. Available from:
- 13. Chong PF, Kira R, Mori H, Okumura A, Torisu H, Yasumoto S, et al. Clinical Features of Acute Flaccid Myelitis Temporally Associated With an Enterovirus D68 Outbreak: Results of a Nationwide Survey of Acute Flaccid Paralysis in Japan, August-December 2015. Clinical infectious diseases: an official publication of the Infectious Diseases Society of America. 2018;66(5):653-64.

DOI:

14. CDC. Acute Flaccid Myelitis - Case Definitions [updated 3 July 2018. Available from:

15. CSTE. Revision to the Standardized Surveillance and Case Definition for Acute Flaccid Myelitis 2017 [updated 2017. Available

from: Agraed and with the control of the control of

16. Ooi MH, Wong SC, Clear D, Perera D, Krishnan S, Preston T, et al. Adenovirus Type 21—Associated Acute Flaccid Paralysis during an Outbreak of Hand-Foot-and-Mouth Disease in Sarawak, Malaysia. Clin Infect Dis. 2003;36(5):550-9.

DOI: AREA TOMPER PLANTED TO

- 17. NDDPH. What is Acute Flaccid Myelitis? [updated November 2017. Available from:
- 18. CDC. About Enterovirus A71 [updated 27 September 2018. Available from:
- 19. Dyda A, Stelzer-Braid S, Adam D, Chughtai AA, MacIntyre CR. The association between acute flaccid myelitis (AFM) and Enterovirus D68 (EV-D68)—what is the evidence for causation? Eurosurveillance. 2018;23(3):17-00310.

DOI:

- 20. Messacar K, Asturias EJ, Hixon AM, Van Leer-Buter C, Niesters HG, Tyler KL, et al. Enterovirus D68 and acute flaccid myelitis—evaluating the evidence for causality. The Lancet Infectious Diseases. 2018. DOI:
- 21. CDC. About Enterovirus D68 [updated 9 October 2018. Available from:
- 22. Royston L, Tapparel C. Rhinoviruses and Respiratory Enteroviruses: Not as Simple as ABC. Viruses. 2016;8(1):16. DOI:
- 23. Knoester M, Helfferich J, Poelman R, Van Leer-Buter C, Brouwer OF, Niesters HGM. Twenty-Nine Cases of Enterovirus-D68 Associated Acute Flaccid Myelitis in Europe 2016; A Case Series and Epidemiologic Overview. Pediatr Infect Dis J. 2018.

- 24. Abad FX, Pinto RM, Bosch A. Survival of enteric viruses on environmental fomites. Appl Environ Microbiol. 1994;60(10):3704-10.
- 25. Shaw J, Welch TR, Milstone AM. The role of syndromic surveillance in directing the public health response to the enterovirus D68 epidemic. JAMA pediatrics.
- 2014;168(11):981-2. DOI:
- 26. Huang Y-JS, Higgs S, Horne KM, Vanlandingham DL. Flavivirus-mosquito interactions. Viruses. 2014;6(11):4703-30. DOI:
- 27. CDC. About Adenoviruses Transmission [updated 23 May 2017. Available from:
- 28. Mills S. 2 In Wisconsin Have Rare Paralysis-Inducing Illness [updated 16 October 2018. Available from:
- 29. Rettner R. More 'polio-like' illnesses reported in Pennsylvania [updated 11 October 2018. Available from:
- 30. Cohen E. Parents accuse CDC of not reporting children's deaths from polio-like AFM [updated 13 November 2018. Available from:
- 31. CDC. Acute Flaccid Myelitis: Interim Considerations for Clinical Management 2014 [updated 7 November 2014. Available from:
- 32. CDC. Saint Louis Encephalitis [updated 31 May 2018. Available from:
- 33. CDC. West Nile virus [updated 30 October 2018. Available from:
- 34. CDC. Japanese Encephalitis Vaccine [updated 5 August 2015. Available from:

- 35. Messacar K, Sillau S, Hopkins SE, Otten C, Wilson-Murphy M, Wong B, et al. Safety, tolerability, and efficacy of fluoxetine as an antiviral for acute flaccid myelitis. Neurology. 2018. DOI:
- 36. Sejvar JJ, Lopez AS, Cortese MM, Leshem E, Pastula DM, Miller L, et al. Acute Flaccid Myelitis in the United States, August–December 2014: Results of Nationwide Surveillance. Clin Infect Dis. 2016;63(6):737-45.

37. Bonwitt J, Poel A, DeBolt C, Gonzales E, Lopez A, Routh J, et al. Acute flaccid myelitis among children—Washington, September—November 2016. MMWR Morbidity and mortality weekly report. 2017;66(31):826.

DOI:

- 38. Sejvar JJ, Lopez AS, Cortese MM, Leshem E, Pastula DM, Miller L, et al. Acute Flaccid Myelitis in the United States, August-December 2014: Results of Nationwide Surveillance. Clin Infect Dis. 2016. DOI:
- 39. Dyda A, Stelzer-Braid S, Adam D, Chughtai AA, MacIntyre CR. The association between acute flaccid myelitis (AFM) and Enterovirus D68 (EV-D68) what is the evidence for causation? Eurosurveillance. 2018;23(3):17-00310.

DOI:

- 40. Hixon AM, Yu G, Leser JS, Yagi S, Clarke P, Chiu CY, et al. A mouse model of paralytic myelitis caused by enterovirus D68. PLoS Pathog. 2017;13(2):e1006199. DOI:
- 41. Antona D, Kossorotoff M, Schuffenecker I, Mirand A, Leruez-Ville M, Bassi C, et al. Severe paediatric conditions linked with EV-A71 and EV-D68, France, May to October 2016. Eurosurveillance. 2016;21(46). DOI:

2

42. Messacar K, Burakoff A, Nix WA, Rogers S, Oberste MS, Gerber SI, et al. Notes from the Field: Enterovirus A71 Neurologic Disease in Children—Colorado, 2018.

Morbidity and Mortality Weekly Report. 2018;67(36):1017.

DOI: The second of the second

- 43. Aliabadi N, Messacar K, Pastula DM, Robinson CC, Leshem E, Sejvar JJ, et al. Enterovirus D68 infection in children with acute flaccid myelitis, Colorado, USA, 2014. Emerg Infect Dis. 2016;22(8):1387. DOI:
- 44. Pons-Salort M, Oberste MS, Pallansch MA, Abedi GR, Takahashi S, Grenfell BT, et al. The seasonality of nonpolio enteroviruses in the United States: Patterns and drivers. Proceedings of the National Academy of Sciences. 2018;115(12):3078.

DOI:

- 45. NYDOH. NYS Department of Health Confirms Cases of Serious Respiratory Virus 2018 [updated 12 October 2018. Available from:
- 46. Pons-Salort M, Grassly NC. Serotype-specific immunity explains the incidence of diseases caused by human enteroviruses. Science. 2018;361(6404):800-3.

DOI:

47. NJDOH. Total of 5 Adenovirus Cases Confirmed at Voorhees Pediatric Facility 2018 [updated 5 November 2018. Available

from:

.=

48. Nedelman M. 30 sickened in adenovirus outbreak in New Jersey, including 10 children who have died: CNN; 2018 [updated 7 November 2018. Available from: